

Miebo[™]
(perfluorohexyloctane
ophthalmic solution)

Patient Profiles

[VIEW PROFILES](#)

INDICATION


MIEBO[®] (perfluorohexyloctane ophthalmic solution) is indicated for the treatment of the signs and symptoms of dry eye disease.

Please see additional Important Safety Information [here](#).
Click [here](#) for full Prescribing Information for MIEBO.

Not actual patient.

MIEBO Patient Profiles

Choose a profile below to get started



Lauren, 39
UNDIAGNOSED,
SYMPTOMATIC



Jack, 36
UNDIAGNOSED,
SYMPTOMATIC



Rebecca, 51
TREATMENT
EXPERIENCED



Not actual patients.

Please see Important Safety Information [here](#).
Click [here](#) for full Prescribing Information for MIEBO.

Miebo[™]
(perfluorohexyloctane
ophthalmic solution)



Is there more to Lauren's eye discomfort?



Meet Lauren, 39

UNDIAGNOSED,
SYMPTOMATIC

“ I've always stayed on top of my health. As soon as my eyes started bothering me, I knew it was time to get them checked out. ”

Not actual patient.

DIAGNOSIS

- Currently undiagnosed with dry eye disease (DED)

MEDICAL HISTORY

- No significant ocular history prior to exam
- Used Accutane® as a teenager

LIFESTYLE FACTORS

- Loves outdoor adventures, like hiking and fishing with her husband and dog
- Works long hours as an ER nurse

CURRENT SYMPTOMS

- Eye dryness that typically improves with blinking; no signs of inflammation

CURRENT/PRIOR TREATMENTS

- Has tried several OTC eye drops with little effect



Lauren doesn't want **dryness** and **constant blinking** to get in the way of the life she loves with her husband and dog—or her work with her patients. After **trying some OTC options**, she's ready to tackle her symptoms head-on.

With symptoms pointing to evaporative DED, consider MIEBO

IMPORTANT SAFETY INFORMATION (CONTINUED)

- MIEBO should not be administered while wearing contact lenses. Contact lenses should be removed before use and for at least 30 minutes after administration of MIEBO

Please see additional Important Safety Information [here](#).
Click [here](#) for full Prescribing Information for MIEBO.

Miebo[™]
(perfluorohexyloctane
ophthalmic solution)



Is there more to Jack's vision changes?

DIAGNOSIS

- Currently undiagnosed with dry eye disease (DED)

MEDICAL HISTORY

- No significant ocular history prior to exam
- Wears glasses

LIFESTYLE FACTORS

- Games online every night with friends
- Works as a freelance web designer

CURRENT SIGNS

- Some recent vision changes motivated Jack to come in and update his glasses prescription (not realizing it could be DED)
- There is evidence of meibomian gland obstruction

CURRENT/PRIOR TREATMENTS

- None



With **both work and play** so **dependent on the quality of his vision**, Jack is eager to address whatever is **causing his evaporative dry eye symptoms**.

With signs pointing to evaporative DED, consider MIEBO

IMPORTANT SAFETY INFORMATION (CONTINUED)

- Instruct patients to instill one drop of MIEBO into each eye four times daily

Please see additional Important Safety Information [here](#).
Click [here](#) for full Prescribing Information for MIEBO.

Miebo[™]
(perfluorohexyloctane
ophthalmic solution)



Meet Jack, 36

**UNDIAGNOSED,
SYMPTOMATIC**

“ Screen time is a big part of my life; whether it's work or gaming, I'm constantly online. Lately, that's started to bother my eyes. ”

Not actual patient.



Meet Rebecca, 51

TREATMENT EXPERIENCED

“ I am a flight attendant, and the dry air irritates my eyes. The last treatment didn't work out for me, so what else is there to try? ”

Not actual patient.

What should Rebecca try next?

DIAGNOSIS

- Diagnosed with dry eye disease (DED) one year ago

MEDICAL HISTORY

- Tried Rx treatment for DED for 6 months but stopped after experiencing side effects and not seeing results
- Is experiencing hormonal changes due to menopause

LIFESTYLE FACTORS

- Enjoys reading
- Works as a flight attendant

CURRENT SYMPTOMS

- Eye dryness, intermittent blurry vision, light sensitivity, and watery eyes

CURRENT/PRIOR TREATMENTS

- Used Rx treatment in the past; now using artificial tears because her symptoms have come back



While Rebecca was proactive in attempting to treat her symptoms, she **isn't getting the results** she needs and might **benefit from a different approach**, preferably one with **minimal side effects**.

With a history of evaporative DED, consider MIEBO

IMPORTANT SAFETY INFORMATION (CONTINUED)

- The safety and efficacy in pediatric patients below the age of 18 have not been established

Please see additional Important Safety Information [here](#).
Click [here](#) for full Prescribing Information for MIEBO.

Miebo[™]
(perfluorohexyloctane
ophthalmic solution)



Indication & Important Safety Information

INDICATION

MIEBO® (perfluorohexyloctane ophthalmic solution) is indicated for the treatment of the signs and symptoms of dry eye disease.

IMPORTANT SAFETY INFORMATION

- MIEBO should not be administered while wearing contact lenses. Contact lenses should be removed before use and for at least 30 minutes after administration of MIEBO
- Instruct patients to instill one drop of MIEBO into each eye four times daily
- The safety and efficacy in pediatric patients below the age of 18 have not been established
- The most common ocular adverse reaction was blurred vision (1% to 3% of patients reported blurred vision and conjunctival redness)

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call [1-800-FDA-1088](tel:1-800-FDA-1088).

Click [here](#) for full Prescribing Information for MIEBO.

BAUSCH + LOMB

Any brand names are trademarks of the respective owners.

©2024 Bausch + Lomb

MBO.0583.USA.23

Miebo[™]
(perfluorohexyloctane
ophthalmic solution)